
Apocalyptic quantum gravity



David Wakeham UBC Gravity Seminar April 28, 2021



Regular gravity is good for understanding

 Regular gravity is good for understanding the universe at large scales


► Regular gravity is good for understanding the universe at large scales and the astrophysical objects within it.

- Regular gravity is good for understanding the universe at large scales and the astrophysical objects within it.
- So is quantum gravity!


- Regular gravity is good for understanding the universe at large scales and the astrophysical objects within it.
- So is quantum gravity! Big Bang

- Regular gravity is good for understanding the universe at large scales and the astrophysical objects within it.
- ► So is quantum gravity! Big Bang and black holes.

- ► Regular gravity is good for understanding the universe at large scales and the astrophysical objects within it.
- ► So is quantum gravity! Big Bang and black holes.




- ► Regular gravity is good for understanding the universe at large scales and the astrophysical objects within it.
- ► So is quantum gravity! Big Bang and black holes.




► The "easy" problem is black holes

- ► Regular gravity is good for understanding the universe at large scales and the astrophysical objects within it.
- ► So is quantum gravity! Big Bang and black holes.



► The "easy" problem is black holes because we can plonk them in a different-looking spacetime,

- ► Regular gravity is good for understanding the universe at large scales and the astrophysical objects within it.
- ► So is quantum gravity! Big Bang and black holes.



► The "easy" problem is black holes because we can plonk them in a different-looking spacetime, AdS/CFT.

Let's consider quantum gravity


Let's consider quantum gravity in negatively curved spacetime, $\Lambda < 0$

Let's consider quantum gravity in negatively curved spacetime, $\Lambda < 0$ (Anti de Sitter space = AdS).

Let's consider quantum gravity in negatively curved spacetime, $\Lambda < 0$ (Anti de Sitter space = AdS).




Let's consider quantum gravity in negatively curved spacetime, $\Lambda < 0$ (Anti de Sitter space = AdS).




Wacky claim:¹

Let's consider quantum gravity in negatively curved spacetime, $\Lambda < 0$ (Anti de Sitter space = AdS).




▶ Wacky claim:¹ this is "dual" to

Let's consider quantum gravity in negatively curved spacetime, $\Lambda < 0$ (Anti de Sitter space = AdS).



 Wacky claim:¹ this is "dual" to a special QFT (conformal field theory = CFT)

Let's consider quantum gravity in negatively curved spacetime, $\Lambda < 0$ (Anti de Sitter space = AdS).



Wacky claim:¹ this is "dual" to a special QFT (conformal field theory = CFT) at the asymptotic boundary.

► A regular QFT has Lorentz invariance

▶ A regular QFT has Lorentz invariance (isometries of $\eta_{\mu\nu}$).


► A regular QFT has Lorentz invariance (isometries of $\eta_{\mu\nu}$). The action is invariant

► A regular QFT has Lorentz invariance (isometries of $\eta_{\mu\nu}$). The action is invariant and fields are covariant.


- ▶ A regular QFT has Lorentz invariance (isometries of $\eta_{\mu\nu}$). The action is invariant and fields are covariant.
- A CFT also has scale invariance,

- ► A regular QFT has Lorentz invariance (isometries of $\eta_{\mu\nu}$). The action is invariant and fields are covariant.
- ▶ A CFT also has scale invariance, i.e. zooming in and out:


- ▶ A regular QFT has Lorentz invariance (isometries of $\eta_{\mu\nu}$). The action is invariant and fields are covariant.
- ▶ A CFT also has scale invariance, i.e. zooming in and out:



- ▶ A regular QFT has Lorentz invariance (isometries of $\eta_{\mu\nu}$). The action is invariant and fields are covariant.
- ▶ A CFT also has scale invariance, i.e. zooming in and out:




- ▶ A regular QFT has Lorentz invariance (isometries of $\eta_{\mu\nu}$). The action is invariant and fields are covariant.
- ▶ A CFT also has scale invariance, i.e. zooming in and out:




► For AdS/CFT, we focus on large-*N* CFTs,

- ▶ A regular QFT has Lorentz invariance (isometries of $\eta_{\mu\nu}$). The action is invariant and fields are covariant.
- ▶ A CFT also has scale invariance, i.e. zooming in and out:



► For AdS/CFT, we focus on large-*N* CFTs, which are almost free.²

- ► A regular QFT has Lorentz invariance (isometries of $\eta_{\mu\nu}$). The action is invariant and fields are covariant.
- ▶ A CFT also has scale invariance, i.e. zooming in and out:



► For AdS/CFT, we focus on large-N CFTs, which are almost free.² Simple fields have vertices suppressed by N.

States and metrics

States and metrics


► Some CFT states correspond to semiclassical geometries

States and metrics


► Some CFT states correspond to semiclassical geometries which "fill in" the cylinder.

► Some CFT states correspond to semiclassical geometries which "fill in" the cylinder. Example: the vacuum state.

► Some CFT states correspond to semiclassical geometries which "fill in" the cylinder. Example: the vacuum state.




▶ Some CFT states correspond to semiclassical geometries which "fill in" the cylinder. Example: the vacuum state.




► More interesting:

▶ Some CFT states correspond to semiclassical geometries which "fill in" the cylinder. Example: the vacuum state.




More interesting: canonical thermal state,

► Some CFT states correspond to semiclassical geometries which "fill in" the cylinder. Example: the vacuum state.




More interesting: canonical thermal state, where the probability of observing $|E\rangle$

▶ Some CFT states correspond to semiclassical geometries which "fill in" the cylinder. Example: the vacuum state.



More interesting: canonical thermal state, where the probability of observing $|E\rangle$ is proportional to $e^{-\beta E}$.

▶ Some CFT states correspond to semiclassical geometries which "fill in" the cylinder. Example: the vacuum state.



- More interesting: canonical thermal state, where the probability of observing $|E\rangle$ is proportional to $e^{-\beta E}$.
- ► This turns out to be dual to an AdS black hole!³

${\sf Eternal/external}$

With this explicit quantum description,


With this explicit quantum description, we might hope to understand

▶ With this explicit quantum description, we might hope to understand (a) the interior,


► With this explicit quantum description, we might hope to understand (a) the interior, and (b) evaporation.⁴

- ► With this explicit quantum description, we might hope to understand (a) the interior, and (b) evaporation.⁴
- ▶ These are good reasons to study quantum gravity!

- ► With this explicit quantum description, we might hope to understand (a) the interior, and (b) evaporation.⁴
- ▶ These are good reasons to study quantum gravity!




- ► With this explicit quantum description, we might hope to understand (a) the interior, and (b) evaporation.⁴
- These are good reasons to study quantum gravity!




▶ Problem:

- ► With this explicit quantum description, we might hope to understand (a) the interior, and (b) evaporation.⁴
- These are good reasons to study quantum gravity!




Problem: we can't do either!

- ► With this explicit quantum description, we might hope to understand (a) the interior, and (b) evaporation.⁴
- ▶ These are good reasons to study quantum gravity!



 Problem: we can't do either! It's just the static exterior geometry,

- ► With this explicit quantum description, we might hope to understand (a) the interior, and (b) evaporation.⁴
- ▶ These are good reasons to study quantum gravity!



▶ Problem: we can't do either! It's just the static exterior geometry, so no interior or evaporation.

► There is no interior


► There is no interior because we have a mixed, i.e. uncertain state.

► There is no interior because we have a mixed, i.e. uncertain state. There is uncertainty about what's inside!


- There is no interior because we have a mixed, i.e. uncertain state. There is uncertainty about what's inside!
- Maybe we can find a pure state

- There is no interior because we have a mixed, i.e. uncertain state. There is uncertainty about what's inside!
- ▶ Maybe we can find a pure state that "looks" thermal.

- There is no interior because we have a mixed, i.e. uncertain state. There is uncertainty about what's inside!
- ▶ Maybe we can find a pure state that "looks" thermal.




- There is no interior because we have a mixed, i.e. uncertain state. There is uncertainty about what's inside!
- Maybe we can find a pure state that "looks" thermal.




On two identical CFTs, the thermofield double (TFD)

- There is no interior because we have a mixed, i.e. uncertain state. There is uncertainty about what's inside!
- ▶ Maybe we can find a pure state that "looks" thermal.



• On two identical CFTs, the thermofield double (TFD) purifies ρ_{β} .

- There is no interior because we have a mixed, i.e. uncertain state. There is uncertainty about what's inside!
- Maybe we can find a pure state that "looks" thermal.



▶ On two identical CFTs, the thermofield double (TFD) purifies ρ_{β} . The dual geometry is the AdS wormhole.⁵

▶ Problem:

▶ Problem: this is a state on two CFTs!

▶ Problem: this is a state on two CFTs! We'd like to get rid of that fictitious second CFT

▶ Problem: this is a state on two CFTs! We'd like to get rid of that fictitious second CFT and still have some interior.

- ▶ Problem: this is a state on two CFTs! We'd like to get rid of that fictitious second CFT and still have some interior.
- Geometrically, could hope to fold the TFD in half.

- ▶ Problem: this is a state on two CFTs! We'd like to get rid of that fictitious second CFT and still have some interior.
- Geometrically, could hope to fold the TFD in half.





- ▶ Problem: this is a state on two CFTs! We'd like to get rid of that fictitious second CFT and still have some interior.
- Geometrically, could hope to fold the TFD in half.





Quantum-mechanically, we can assure a pure state

A folding projector

- ▶ Problem: this is a state on two CFTs! We'd like to get rid of that fictitious second CFT and still have some interior.
- Geometrically, could hope to fold the TFD in half.





 Quantum-mechanically, we can assure a pure state by projecting the second CFT

A folding projector

- ▶ Problem: this is a state on two CFTs! We'd like to get rid of that fictitious second CFT and still have some interior.
- Geometrically, could hope to fold the TFD in half.





▶ Quantum-mechanically, we can assure a pure state by projecting the second CFT onto some fixed state |B⟩:

A folding projector

- Problem: this is a state on two CFTs! We'd like to get rid of that fictitious second CFT and still have some interior.
- Geometrically, could hope to fold the TFD in half.





▶ Quantum-mechanically, we can assure a pure state by projecting the second CFT onto some fixed state |B⟩:

$$|\mathsf{TFD}_{\beta}\rangle_{12} \to \langle B|\mathsf{TFD}_{\beta}\rangle_{1} = |B_{\beta}\rangle_{1}.$$




▶ It's not at all obvious how to do this!


It's not at all obvious how to do this! So let's do a magic trick,

► It's not at all obvious how to do this! So let's do a magic trick, and Wick rotate real to imaginary time,


▶ It's not at all obvious how to do this! So let's do a magic trick, and Wick rotate real to imaginary time, $\tau = it$:

▶ It's not at all obvious how to do this! So let's do a magic trick, and Wick rotate real to imaginary time, $\tau = it$:







▶ It's not at all obvious how to do this! So let's do a magic trick, and Wick rotate real to imaginary time, $\tau = it$:



▶ This fuses the hyperbola to a circle,


▶ It's not at all obvious how to do this! So let's do a magic trick, and Wick rotate real to imaginary time, $\tau = it$:






▶ This fuses the hyperbola to a circle, and two CFTs to one!

▶ It's not at all obvious how to do this! So let's do a magic trick, and Wick rotate real to imaginary time, $\tau = it$:



- ▶ This fuses the hyperbola to a circle, and two CFTs to one!
- ▶ The CFT is in a thermal state,

▶ It's not at all obvious how to do this! So let's do a magic trick, and Wick rotate real to imaginary time, $\tau = it$:




- ► This fuses the hyperbola to a circle, and two CFTs to one!
- ► The CFT is in a thermal state, geometrized by a "thermal" circle on the outside.

▶ On the CFT, consider states which are mirror symmetric.


- ▶ On the CFT, consider states which are mirror symmetric.
- ▶ On the circle, we can evolve to the top and bottom!

- ▶ On the CFT, consider states which are mirror symmetric.
- ► On the circle, we can evolve to the top and bottom! This gives us states which are "local" in time.

- ▶ On the CFT, consider states which are mirror symmetric.
- ► On the circle, we can evolve to the top and bottom! This gives us states which are "local" in time.




- ▶ On the CFT, consider states which are mirror symmetric.
- ► On the circle, we can evolve to the top and bottom! This gives us states which are "local" in time.




► These special "mirror" states are called boundary states.

- ▶ On the CFT, consider states which are mirror symmetric.
- ► On the circle, we can evolve to the top and bottom! This gives us states which are "local" in time.



► These special "mirror" states are called boundary states. They cut the outer circle in half,

- On the CFT, consider states which are mirror symmetric.
- ► On the circle, we can evolve to the top and bottom! This gives us states which are "local" in time.



► These special "mirror" states are called boundary states. They cut the outer circle in half, but what happens inside?


► Boundary states break imaginary time translation.

- ► Boundary states break imaginary time translation.
- We can foliate spacetime


- ► Boundary states break imaginary time translation.
- ► We can foliate spacetime with hypersurfaces satisfying the remaining symmetries

- ► Boundary states break imaginary time translation.
- ▶ We can foliate spacetime with hypersurfaces satisfying the remaining symmetries (anchored at the top and bottom).

- ► Boundary states break imaginary time translation.
- ▶ We can foliate spacetime with hypersurfaces satisfying the remaining symmetries (anchored at the top and bottom).




- ► Boundary states break imaginary time translation.
- ▶ We can foliate spacetime with hypersurfaces satisfying the remaining symmetries (anchored at the top and bottom).



These hypersurface are the fabled end-of-the-world (EOW) branes.

- Boundary states break imaginary time translation.
- ▶ We can foliate spacetime with hypersurfaces satisfying the remaining symmetries (anchored at the top and bottom).



These hypersurface are the fabled end-of-the-world (EOW) branes. Our spacetime could end on any of them!


► To return to real time,

► To return to real time, we glue real and imaginary diagrams along a horizontal slice.


- ➤ To return to real time, we glue real and imaginary diagrams along a horizontal slice.
- Branes continue at fixed tension

- ➤ To return to real time, we glue real and imaginary diagrams along a horizontal slice.
- ▶ Branes continue at fixed tension (cf. symmetry).

- ➤ To return to real time, we glue real and imaginary diagrams along a horizontal slice.
- ▶ Branes continue at fixed tension (cf. symmetry).




- ► To return to real time, we glue real and imaginary diagrams along a horizontal slice.
- ▶ Branes continue at fixed tension (cf. symmetry).



We are still folding spacetime in half,

Back to reality


- ► To return to real time, we glue real and imaginary diagrams along a horizontal slice.
- ▶ Branes continue at fixed tension (cf. symmetry).



➤ We are still folding spacetime in half, but it's "distended".

Back to reality

- ► To return to real time, we glue real and imaginary diagrams along a horizontal slice.
- Branes continue at fixed tension (cf. symmetry).



We are still folding spacetime in half, but it's "distended". Boundary states modify the bulk!


• It seems the pure states $|B_{\beta}\rangle = \langle B|\mathsf{TFD}_{\beta}\rangle$

▶ It seems the pure states $|B_{\beta}\rangle = \langle B|\mathsf{TFD}_{\beta}\rangle$ describe black holes with some interior.


▶ It seems the pure states $|B_{\beta}\rangle = \langle B|\mathsf{TFD}_{\beta}\rangle$ describe black holes with some interior. But what can we do with this?

- ▶ It seems the pure states $|B_{\beta}\rangle = \langle B|\mathsf{TFD}_{\beta}\rangle$ describe black holes with some interior. But what can we do with this?
- An important tool is entanglement!

- ▶ It seems the pure states $|B_{\beta}\rangle = \langle B|\mathsf{TFD}_{\beta}\rangle$ describe black holes with some interior. But what can we do with this?
- An important tool is entanglement!




- ▶ It seems the pure states $|B_{\beta}\rangle = \langle B|\mathsf{TFD}_{\beta}\rangle$ describe black holes with some interior. But what can we do with this?
- An important tool is entanglement!




► Loosely speaking, systems A and B are entangled

- ▶ It seems the pure states $|B_{\beta}\rangle = \langle B|\mathsf{TFD}_{\beta}\rangle$ describe black holes with some interior. But what can we do with this?
- An important tool is entanglement!



► Loosely speaking, systems *A* and *B* are entangled when they share EPR pairs.

- ▶ It seems the pure states $|B_{\beta}\rangle = \langle B|\mathsf{TFD}_{\beta}\rangle$ describe black holes with some interior. But what can we do with this?
- An important tool is entanglement!



▶ Loosely speaking, systems *A* and *B* are entangled when they share EPR pairs. The systems above share three.

• Given a density matrix ρ_A for system A,


• Given a density matrix ρ_A for system A, uncertainty can come from

▶ Given a density matrix ρ_A for system A, uncertainty can come from (a) entanglement

▶ Given a density matrix ρ_A for system A, uncertainty can come from (a) entanglement and (b) classical entropy.


- Given a density matrix ρ_A for system A, uncertainty can come from (a) entanglement and (b) classical entropy.
- ▶ Either way, we have uncertainty about the state on A.

- Given a density matrix ρ_A for system A, uncertainty can come from (a) entanglement and (b) classical entropy.
- ▶ Either way, we have uncertainty about the state on *A*.




$$S[
ho_A] = - \left\langle \log
ho_A \right
angle_{
ho_A}$$

- Given a density matrix ρ_A for system A, uncertainty can come from (a) entanglement and (b) classical entropy.
- ▶ Either way, we have uncertainty about the state on *A*.



▶ A good measure is the entanglement entropy⁶ $S[\rho_A]$.

- ▶ Given a density matrix ρ_A for system A, uncertainty can come from (a) entanglement and (b) classical entropy.
- ▶ Either way, we have uncertainty about the state on *A*.




A good measure is the entanglement entropy⁶ $S[\rho_A]$. (It's just the Shannon entropy for the distribution over states.)

▶ In AdS/CFT, we'll look at spatial subsystems A.


► In AdS/CFT, we'll look at spatial subsystems A. In states with a nice gravity dual, a miracle occurs:⁷

► In AdS/CFT, we'll look at spatial subsystems A. In states with a nice gravity dual, a miracle occurs:⁷



$$S[
ho_A] = \min_{\gamma_A} rac{ ext{Area}[\gamma_A]}{4\,G_{ ext{N}}}$$


► In AdS/CFT, we'll look at spatial subsystems A. In states with a nice gravity dual, a miracle occurs:⁷



$$extit{S}[
ho_A] = \min_{\gamma_A} rac{ ext{Area}[\gamma_A]}{4\,G_{ ext{N}}}$$

Ryu-Takayanagi (RT):


► In AdS/CFT, we'll look at spatial subsystems A. In states with a nice gravity dual, a miracle occurs:⁷



$$extit{S}[
ho_{ extit{A}}] = \min_{\gamma_{ extit{A}}} rac{ ext{Area}[\gamma_{ extit{A}}]}{4\,G_{ ext{N}}}$$

ightharpoonup Ryu-Takayanagi (RT): find the minimal area surface γ_A


► In AdS/CFT, we'll look at spatial subsystems A. In states with a nice gravity dual, a miracle occurs:⁷



$$S[
ho_A] = \min_{\gamma_A} rac{ ext{Area}[\gamma_A]}{4\,G_{ ext{N}}}$$

▶ Ryu-Takayanagi (RT): find the minimal area surface γ_A which is anchored at A,


► In AdS/CFT, we'll look at spatial subsystems A. In states with a nice gravity dual, a miracle occurs:⁷



$$S[
ho_A] = \min_{\gamma_A} rac{ ext{Area}[\gamma_A]}{4\,G_{ ext{N}}}$$

▶ Ryu-Takayanagi (RT): find the minimal area surface γ_A which is anchored at A, and divide its area by $4G_N \sim 1/N$.


► In AdS/CFT, we'll look at spatial subsystems A. In states with a nice gravity dual, a miracle occurs:⁷



$$S[
ho_A] = \min_{\gamma_A} rac{ ext{Area}[\gamma_A]}{4\,G_{ ext{N}}}$$

- ▶ Ryu-Takayanagi (RT): find the minimal area surface γ_A which is anchored at A, and divide its area by $4G_N \sim 1/N$.
- ► The purple entanglement wedge

► In AdS/CFT, we'll look at spatial subsystems A. In states with a nice gravity dual, a miracle occurs:⁷



$$S[
ho_A] = \min_{\gamma_A} rac{ ext{Area}[\gamma_A]}{4\,G_{ ext{N}}}$$


- ▶ Ryu-Takayanagi (RT): find the minimal area surface γ_A which is anchored at A, and divide its area by $4G_N \sim 1/N$.
- ▶ The purple entanglement wedge is encoded by ρ_A .8


▶ What happens with an EOW brane?

▶ What happens with an EOW brane? Conjecture:


▶ What happens with an EOW brane? Conjecture: the minimal surfaces γ_A can end on the brane.


▶ What happens with an EOW brane? Conjecture: the minimal surfaces γ_A can end on the brane.





▶ What happens with an EOW brane? Conjecture: the minimal surfaces γ_A can end on the brane.







► Since branes are like mirrors,

Apocalyptic RT

▶ What happens with an EOW brane? Conjecture: the minimal surfaces γ_A can end on the brane.







► Since branes are like mirrors, when we compute entanglement entropy for *A*,

Apocalyptic RT

▶ What happens with an EOW brane? Conjecture: the minimal surfaces γ_A can end on the brane.







▶ Since branes are like mirrors, when we compute entanglement entropy for A, we really do it for $A \cup A_{mirror}$.

Apocalyptic RT

▶ What happens with an EOW brane? Conjecture: the minimal surfaces γ_A can end on the brane.






- ▶ Since branes are like mirrors, when we compute entanglement entropy for A, we really do it for $A \cup A_{\text{mirror}}$.
- ► (It's the method of images for quantum gravity.)


Let's use our apocalyptic RT formula in the black hole.

- Let's use our apocalyptic RT formula in the black hole.
- ▶ Since ρ_A is dual to the purple entanglement wedge,


- ▶ Let's use our apocalyptic RT formula in the black hole.
- Since ρ_A is dual to the purple entanglement wedge, we should have access to physics behind the horizon.

- ▶ Let's use our apocalyptic RT formula in the black hole.
- ▶ Since ρ_A is dual to the purple entanglement wedge, we should have access to physics behind the horizon.






- ▶ Let's use our apocalyptic RT formula in the black hole.
- ▶ Since ρ_A is dual to the purple entanglement wedge, we should have access to physics behind the horizon.




▶ We did explicit calculations to find regimes γ_A falls in.

- Let's use our apocalyptic RT formula in the black hole.
- ▶ Since ρ_A is dual to the purple entanglement wedge, we should have access to physics behind the horizon.




▶ We did explicit calculations to find regimes γ_A falls in. You need large A

- Let's use our apocalyptic RT formula in the black hole.
- ▶ Since ρ_A is dual to the purple entanglement wedge, we should have access to physics behind the horizon.




We did explicit calculations⁹ to find regimes γ_A falls in. You need large A and a close enough brane to hit it.




▶ We have an elegant semiclassical recipe for a black hole interior.

We have an elegant semiclassical recipe for a black hole interior. But what stops something going terribly wrong?

We have an elegant semiclassical recipe for a black hole interior. But what stops something going terribly wrong?




We have an elegant semiclassical recipe for a black hole interior. But what stops something going terribly wrong?




Spacetime may not end sharply,

We have an elegant semiclassical recipe for a black hole interior. But what stops something going terribly wrong?




Spacetime may not end sharply, or even exist!

We have an elegant semiclassical recipe for a black hole interior. But what stops something going terribly wrong?



- Spacetime may not end sharply, or even exist!
- We need to descend into the microscopic realm

We have an elegant semiclassical recipe for a black hole interior. But what stops something going terribly wrong?



- Spacetime may not end sharply, or even exist!
- We need to descend into the microscopic realm and perform some explicit checks.


► The most tractable setting is a 2d CFT,

► The most tractable setting is a 2d CFT, dual to 3d gravity.


► The most tractable setting is a 2d CFT, dual to 3d gravity. Minimal surfaces are just geodesics!

- ► The most tractable setting is a 2d CFT, dual to 3d gravity. Minimal surfaces are just geodesics!
- ▶ We can also compute S_A on the CFT side: ¹⁰

- ► The most tractable setting is a 2d CFT, dual to 3d gravity. Minimal surfaces are just geodesics!
- ▶ We can also compute S_A on the CFT side: 10




- ► The most tractable setting is a 2d CFT, dual to 3d gravity. Minimal surfaces are just geodesics!
- ▶ We can also compute S_A on the CFT side: 10




► This agrees with the gravity calculation for large central charge *c*

- ► The most tractable setting is a 2d CFT, dual to 3d gravity. Minimal surfaces are just geodesics!
- ▶ We can also compute S_A on the CFT side: 10



► This agrees with the gravity calculation for large central charge c (the equivalent of N)

- ► The most tractable setting is a 2d CFT, dual to 3d gravity. Minimal surfaces are just geodesics!
- ▶ We can also compute S_A on the CFT side: 10



▶ This agrees with the gravity calculation for large central charge c (the equivalent of N) and excitations $O(c^0)$.


▶ We need to find a CFT parameter which bends the brane.

- ▶ We need to find a CFT parameter which bends the brane.
- ▶ The relevant parameter is g,


- ▶ We need to find a CFT parameter which bends the brane.
- ► The relevant parameter is g, the boundary entropy, 11

- We need to find a CFT parameter which bends the brane.
- The relevant parameter is g, the boundary entropy, ¹¹ most simply defined as the vacuum overlap $g = \langle B|0\rangle$.

- We need to find a CFT parameter which bends the brane.
- The relevant parameter is g, the boundary entropy, ¹¹ most simply defined as the vacuum overlap $g = \langle B|0\rangle$.




- We need to find a CFT parameter which bends the brane.
- The relevant parameter is g, the boundary entropy, ¹¹ most simply defined as the vacuum overlap $g = \langle B|0\rangle$.



▶ This validates the apocalyptic RT formula

- We need to find a CFT parameter which bends the brane.
- The relevant parameter is g, the boundary entropy, ¹¹ most simply defined as the vacuum overlap $g = \langle B|0\rangle$.



➤ This validates the apocalyptic RT formula and gives us a microscopic handle on the black hole interior.


► Can we now do stuff behind the horizon?


► Can we now do stuff behind the horizon? Folkore states that ρ_A

► Can we now do stuff behind the horizon? Folkore states that ρ_A "encodes" the entanglement wedge.


- ▶ Can we now do stuff behind the horizon? Folkore states that ρ_A "encodes" the entanglement wedge.
- ▶ So, how do I extract a behind-the-horizon spin from ρ_A ?

- ▶ Can we now do stuff behind the horizon? Folkore states that ρ_A "encodes" the entanglement wedge.
- ▶ So, how do I extract a behind-the-horizon spin from ρ_A ?






- ► Can we now do stuff behind the horizon? Folkore states that ρ_A "encodes" the entanglement wedge.
- ▶ So, how do I extract a behind-the-horizon spin from ρ_A ?




▶ Some aspects have been worked out, 12

- ► Can we now do stuff behind the horizon? Folkore states that ρ_A "encodes" the entanglement wedge.
- ▶ So, how do I extract a behind-the-horizon spin from ρ_A ?



► Some aspects have been worked out, 12 but local recovery with branes remains open.

- ► Can we now do stuff behind the horizon? Folkore states that ρ_A "encodes" the entanglement wedge.
- ▶ So, how do I extract a behind-the-horizon spin from ρ_A ?




► Some aspects have been worked out, 12 but local recovery with branes remains open. Work in progress!


Black hole evaporation is a whole other talk!

▶ Black hole evaporation is a whole other talk! But you can jerry-rig these models to become time-dependent.¹³

▶ Black hole evaporation is a whole other talk! But you can jerry-rig these models to become time-dependent. 13




▶ Black hole evaporation is a whole other talk! But you can jerry-rig these models to become time-dependent. 13




Circling back round to our initial motivation,

▶ Black hole evaporation is a whole other talk! But you can jerry-rig these models to become time-dependent.¹³



 Circling back round to our initial motivation, the brane itself looks like a Big Bang/Big Crunch cosmology.

▶ Black hole evaporation is a whole other talk! But you can jerry-rig these models to become time-dependent. 13



- Circling back round to our initial motivation, the brane itself looks like a Big Bang/Big Crunch cosmology.
- ► The trick is to localize gravity to the brane. 14

Conclusion

Conclusion



Conclusion



Thanks for listening!

References

- Maldacena 97; Witten 98; Gubser, Klebanov, Polyakov 98.
- 't Hooft 74.
- 3. Witten 98.
- 4. Hawking 75.
- 5. Israel 76; Maldacena 01.
- 6. Von Neumann 36; Shannon 48.
- 7. Ryu, Takayanagi 06; Hubeny, Rangamani, Takayanagi 07.
- 8. Czech, Karczmarek, Nogueira, Van Raamsdonk 12; Dong, Harlow, Wall 16.
- 9. Cooper, Rozali, Swingle, Van Raamsdonk, Waddell, DW 18.
- Hartman 13; Sully, Van Raamsdonk, DW 20.
- 11. Affleck, Ludwig 91; Takayanagi 12.
- Almheiri 18.
- 13. Rozali, Sully, Van Raamsdonk, Waddell, Wakeham 19.
- 14. Antonini, Swingle 19.